



# Geometric Modelling Summer 2018

Prof. Dr. Hans Hagen

<http://hci.uni-kl.de/teaching/geometric-modelling-ss2018>



# Affine Spaces, Elliptic and Hyperbolic Geometry



## Affine Spaces

Euclidean Geometry can be formulated by the invariant theories.

**Question:** What is invariant under Euclidean transformations (rotation, translation)?

**Answer:** length of vectors, angles between vectors, volumes spanned by vectors

**Question:** What is invariant under projective transformations?

**Answer:** dimension of subspaces (lines are mapped to lines etc.), cross ratio

In this chapter, we will discuss affine spaces and transformations as well as elliptic and hyperbolic geometry.



## Affine Spaces

### Definition: Affine Space

A non-empty set  $A$  of elements ("points") is called an **Affine Space**, if there exist a vector space  $V$  and a map  $\mapsto$  s.t. every pair  $(p, q)$  of points with  $p, q \in A$  are mapped to exactly one vector  $\vec{v} \in V$  s.t.

- a) For every  $\vec{v} \in V$  and  $p \in A$  there is exactly one point  $q \in A$  with  $(p, q) \mapsto \vec{v}$ .
- b) If  $(p, q) \mapsto \vec{v}$  and  $(q, r) \mapsto \vec{w}$ , then  $(p, r) \mapsto \vec{v} + \vec{w}$ .

$V$  is called the vector space belonging to  $A$ .

A **base of  $A$**  is given by  $A : \{0, \vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$ , where  $0 \in A$  and  $\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}$  form a base of  $V$ .



## Affine Spaces

### Definition: Affine Transformation

A map  $F : A \rightarrow B$  where  $A$  and  $B$  are affine spaces is called an **affine transformation**, if it preserves the colinearity of points and the ratios of vectors along a line.

A **coordinate representation** of such a transformation is given by "a linear transformation  $+ \vec{a}$ ":

$$\vec{y} = A\vec{x} + \vec{a} \quad \text{where } \det(A) \neq 0$$



## Affine Spaces

### properties of affine transformations:

- preserve collinearity and coplanarity of points (i.e.: lines and planes are preserved)
- preserve ratios of vectors along lines
- preserve parallelity



## Affine Spaces

### Definition: Affine Group

Let  $A_n$  denote an  $n$ -dim. affine space. Then,  $\mathcal{A}_n := \{F \mid F : A_n \mapsto A_n, F \text{ nondegenerate (i.e. invertible) and affine}\}$  is a group under composition.  $\mathcal{A}_n$  is called the **affine group**.

### Theorem

The set of all  $n$ -row quadratic matrices whose determinant does not vanish constitutes a group under matrix multiplication. If  $R$  is the scalar field, this group is denoted  $GL(n, R)$ . The real affine group is isomorphic to  $GL(n, \mathbb{R})$ .



## Erlangen Program

### **Felix Klein's Erlangen Program**

Felix Klein proposed to employ groups for a unified and clear treatment of geometry. In this view, the geometry belonging to a group of transformations is the theory of invariants of this group. Therefore, the task of affine geometry is to investigate properties that are invariant under affine transformations.

From group theory, "hierarchies of geometry" arise.





## Erlangen Program

The **Euclidean motions** are special cases of affine transformations that are not only line-, ratio- and parallell-preserving but also preserve metric quantities (angles, lengths, volumes,...). The **Euclidean group** is a subgroup of the affine group.

### Coordinate Representation of the Euclidean Motions

$$\vec{y} = B\vec{x} + \vec{a} \quad \text{where } B \text{ is an orthogonal matrix (i.e.) } B^T = B^{-1}$$

$\det(B) > 0 \rightarrow$  orientation preserving, **direct motions**, rigid motions (examples: rotations, translations)

$\det(B) < 0 \rightarrow$  not preserving the orientation, **indirect motions** (example: axis reflections, point reflections in more than 2d)



## Erlangen Program

### Integration of the Classical Geometries in the Sense of the Erlangen Program:

Consider the subgroup of the projective group (i.e. the corresponding geometry) which fixes a hyper plane. We so to say "tag" this **hyperplane as absolute figure at infinity** and restrict the effect of the subgroup of the transformation group to the points that are not incident with this hyperplane.

This procedure yields affine transformations by restricting the projective transformations to the points that are not at infinity.

The **affine group** then reveals itself as a **subgroup of the projective group**.



## Erlangen Program

"The Procedure in Coordinates":  $P^3$  ( $x_0 = 0$  is mapped to  $y = 0$ )

$$\begin{bmatrix} y_0 \\ y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} a_{00} & 0 & 0 & 0 \\ a_{10} & a_{11} & a_{12} & a_{13} \\ a_{20} & a_{21} & a_{22} & a_{23} \\ a_{30} & a_{31} & a_{32} & a_{33} \end{bmatrix} \cdot \begin{bmatrix} x_0 \\ x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$\rightarrow \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = \begin{pmatrix} a_{10} \cdot x_0 & a_{11} \cdot x_1 & a_{12} \cdot x_2 & a_{13} \cdot x_3 \\ a_{20} \cdot x_0 & a_{21} \cdot x_1 & a_{22} \cdot x_2 & a_{23} \cdot x_3 \\ a_{30} \cdot x_0 & a_{31} \cdot x_1 & a_{32} \cdot x_2 & a_{33} \cdot x_3 \end{pmatrix}$$

→ restrict to the points not at infinity (divide by  $x_0$ )

$$\rightarrow \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \cdot \begin{pmatrix} \frac{x_1}{x_0} & \frac{x_2}{x_0} & \frac{x_3}{x_0} \end{pmatrix}^T + \begin{pmatrix} a_{10} \\ a_{20} \\ a_{30} \end{pmatrix}$$

affine transformation

## Erlangen Program

Not only the affine (and therefore the Euclidean) geometries can be "integrated" into projective geometry. This proposition does also hold for the hyperbolic and elliptic geometries. Unfortunately we cannot discuss the axiomatic structure of geometry thoroughly in this course.


The common axiomatic foundation of the Euclidean, the elliptic, and the hyperbolic geometries is the so-called *absolute geometry*. To this fundamental axiomatic system, one of the interchangeable parallel axioms is added:

- elliptical parallel axiom: There is no parallel line.
- Euclidean parallel axiom: There is exactly one parallel line.
- hyperbolic parallel axiom: There are at least two parallel lines.



## Elliptic and Hyperbolic Geometry


### Model of the Hyperbolic Plane (Beltrami-Klein model):



Points:  $\{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 1\}$

Lines: {secants}

### Model of the Elliptic Plane:



Points:  $\{(P', P'') \mid P, P'' \in S^2 \wedge x'_i = -x''_i\}$

Lines: {great circles}



## Elliptic and Hyperbolic Geometry

**Important Note:** Elliptic and Hyperbolic spaces are metric spaces, i.e. lengths, angles, volumes,... are computable in the sense of elliptical or hyperbolic metrics!



## Elliptic and Hyperbolic Geometry

In later chapters, we will discuss the theory of geodesics. This will enable us to measure the lengths of lines on hyperplanes of arbitrary curvature. As elliptic space have a constant positive and hyperbolic spaces have a constant negative curvature, this will also enable us to study lengths in these geometries. Thus, we will not provide proper equations here.

We will discuss the measurement of angles in hyperbolic spaces, though.


Measuring angles in the Beltrami-Klein model is rather difficult, so we first introduce another model of hyperbolic geometry, the Poincaré disk model. After that, we introduce the hyperbolic functions and discuss triangles in the hyperbolic plane.



## Elliptic and Hyperbolic Geometry

### Poincaré disk model

The Beltrami-Klein model is only conformal in the offspring making it hard to measure angles as their values depend on the position. In the Poincaré disk model, angles are measured like in Euclidean space. The angles between two lines in the Poincaré disk model are the angles between the tangents at the intersection of the two lines.



Points:  $\{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 1\}$

Lines: {arcs}

**Figure:** Two lines (red) and (blue), and the angle between them in the Poincaré disk model.






## Elliptic and Hyperbolic Geometry

### Hyperbolic Functions

The hyperbolic sine and hyperbolic cosine describe the right half of the unit hyperbola  $x^2 - y^2 = 1$  just like sine and cosine describe the unit circle. Moreover, the hyperbolic cosine describes the form of a rope hung up at its two end points.





## Elliptic and Hyperbolic Geometry


The hyperbolic functions are given by the following equations:

### Definition: Hyperbolic Sine, Hyperbolic Cosine, Hyperbolic Tangent

The hyperbolic sine and hyperbolic cosine are the odd and even components of the exponential function:

- **Hyperbolic Sine:**  $\sinh x = \frac{1}{2} (e^x - e^{-x}) = -i \sin(ix)$
- **Hyperbolic Cosine:**  $\cosh x = \frac{1}{2} (e^x + e^{-x}) = \cos(ix)$
- **Hyperbolic Tangent:**  $\tanh x = \frac{\sinh x}{\cosh x}$

## Elliptic and Hyperbolic Geometry



**Figure:** Hyperbolic Sine (red), Hyperbolic Cosine (green), and Hyperbolic Tangent (blue).



## Elliptic and Hyperbolic Geometry

### properties:

- 1  $\frac{d}{dx} \sinh x = \cosh x$ ;  $\frac{d}{dx} \cosh x = \sinh x$
- 2  $\cosh^2 x - \sinh^2 x = 1$
- 3 Euler's Formula:  $\cosh x + \sinh x = e^x$
- 4  $\cosh x - \sinh x = e^{-x}$
- 5 A homogeneous rope that is hung up at its two end points and sags only due to its own weight can be described by a cosh-curve. Such a curve is called a **catenary**.
- 6 The area under the curve of  $\cosh x$  is equal to its arc length:


$$A = \int_a^b \cosh(x) dx = \int_a^b \sqrt{1 + \left(\frac{d}{dx} \cosh(x)\right)^2} dx = \text{arc length}$$



## Elliptic and Hyperbolic Geometry

### Triangle in Hyperbolic Space

As in Euclidean space, a triangle in hyperbolic space is uniquely determined by the intersection of three lines:



**Figure:** Triangle in the Poincaré disk model of the hyperbolic plane.

The sum of the angles in a triangle in hyperbolic spaces is always smaller than  $\pi$ , in elliptic spaces, it is always larger.



## Elliptic and Hyperbolic Geometry

In hyperbolic space, the following theorems similar to the sine rule and the cosine rule in trigonometry hold:

### Theorem

Let  $\alpha$ ,  $\beta$ , and  $\gamma$ , denote the angles of a triangle in hyperbolic space and  $a$ ,  $b$ , and  $c$  the lengths of the opposite sides. Then:

$$\frac{\sin \alpha}{\sinh a} = \frac{\sin \beta}{\sinh b} = \frac{\sin \gamma}{\sinh c}$$

$$\cosh c = \cosh a \cdot \cosh b - \sinh a \cdot \sinh b \cdot \sin \gamma$$



## Elliptic and Hyperbolic Geometry

**Integration into Projective Geometry** We have been able to integrate affine geometry into projective geometry by the distinction of a hyperplane as absolute figure. Analogously, we can integrate the elliptic and hyperbolic geometry by the distinction of special quadrics:

### Definition: Quadric

Let  $f : V \times V \rightarrow K$  be a bilinear form and its corresponding quadratic form  $F = V \rightarrow K; \vec{x} \mapsto f(\vec{x}, \vec{x})$ .

Then,  $Q := \{P(\vec{x}) \in P(V) \mid F(\vec{x}) = 0\}$  is called a 2nd-order hyperplane or **Quadric**.



## Elliptic and Hyperbolic Geometry

### Hyperbolic Geometry:

**absolute figure:**  $x_0^2 + x_1^2 + \dots + x_n^2 - x_{n+1}^2 = 0$

Quadric  $Q$  of rank  $n + 1$  and signature  $(n; 1)$ .

The points where  $x_0^2 + x_1^2 + \dots + x_n^2 < x_{n+1}^2$  are called the quadric's **inner points**.

The group of those projective transformations that fix the quadric and map inner points to inner points constitutes a subgroup of the projective group. This subgroup is isomorphic to the group of hyperbolic motions.





## Elliptic and Hyperbolic Geometry

### Elliptic Geometry:

**absolute figure:**  $x_0^2 + x_1^2 + \dots + x_n^2 + x_{n+1}^2 = 0$

Quadric  $Q$  without real points.

Elliptic geometry supports a projective incidence structure.

The group of those projective transformations that fix the absolute figure is a subgroup of the projective group. This subgroup is isomorphic to the group of elliptic motions.